



---

## IMMUNOGENETICS OF LUPUS ERYTHEMATOSUS

Ashraf Antar El ghzaly

*Clinical pathology department, Faculty of Medicine, Mansoura University, 24 El Gomhoria Street, Mansoura 35516, Egypt*

Shereen Salah Metwally

*Clinical pathology department, Faculty of Medicine, Mansoura University, 24 El Gomhoria Street, Mansoura 35516, Egypt*

Farha El-Chennawi

*Clinical pathology department, Faculty of Medicine, Mansoura University, 24 El Gomhoria Street, Mansoura 35516, Egypt*

Maha Elgayar

*\*Dermatology & Andrology department, Faculty of Medicine, Mansoura University, 24 El Gomhoria Street, Mansoura 35516, Egypt.*

Saleh Mohamad Ibrahim

*Clinic for Dermatology. Lübeck University, 160 Ratzeburger Allee, Lübeck 23562, Germany.*

Follow this and additional works at: <https://mmj.mans.edu.eg/home>

---

### Recommended Citation

El ghzaly, Ashraf Antar; Metwally, Shereen Salah; El-Chennawi, Farha; Elgayar, Maha; and Ibrahim, Saleh Mohamad (2013) "IMMUNOGENETICS OF LUPUS ERYTHEMATOSUS," *Mansoura Medical Journal*: Vol. 42 : Iss. 2 , Article 2. Available at: <https://doi.org/10.21608/mjmu.2020.124907>

This Original Study is brought to you for free and open access by Mansoura Medical Journal. It has been accepted for inclusion in Mansoura Medical Journal by an authorized editor of Mansoura Medical Journal. For more information, please contact [mmj@mans.edu.eg](mailto:mmj@mans.edu.eg).

# IMMUNOGENETICS OF LUPUS ERYTHEMATOSUS

*By*

Ashraf Antar El ghzaly \*, Shereen Salah Metwally \*,  
Farha Abd Elaziz El-Chennawi\*, Maha Abd Elghaffar  
Elgayar\*\* , Saleh Mohamad Ibrahim\*\*\*

*From*

\*Clinical pathology department, Faculty of Medicine, Mansoura University, 24  
El Gomhoria Street, Mansoura 35516, Egypt.

\*\*Dermatology & Andrology department ,Faculty of Medicine,  
Mansoura University, 24 El Gomhoria Street, Mansoura 35516, Egypt.

\*\*\*Clinic for Dermatology. Lübeck University, 160 Ratzeburger Allee,  
Lübeck 23562, Germany.

## ABSTRACT

*Key words* : Lupus , Genetics, Association.

### Immunopathogenesis of SLE

Systemic Lupus Erythematosus (SLE) is a systemic autoimmune disease with a prevalence of approximately 1/2500 in European populations affecting mainly females (9 females:1 male) (1). It's characterized by a plethora of immunological and laboratory abnormalities with multiple organ / tissue damage. Almost all of the key components in

the immune system are involved in the disease pathogenesis which is largely still unclear (2) .

Deficiencies in the recognition and clearance of apoptotic cells by phagocytosis have been shown in patients with SLE (3,4). Whether apoptosis itself is abnormal or merely an effect of environmental triggers, such as U.V. irradiation or viral infection, is less understood (5,6) .

During necrosis or apoptosis nuclear antigens are subjected to modifications which allow them to be rec-

ognized as non-self. Activation of TLR7 and TLR9 by self nucleic acids and immune complexes leads to the production of IFN- $\alpha$  and proinflammatory cytokines such as IL-6, TNF- $\alpha$  and IL-12 (7).

IFN- $\alpha$  fosters autoimmunity by several mechanisms; It promotes dendritic cell maturation, the production of pro-inflammatory cytokines, stimulation of Th1 pathways, B-cell activation, plasma cell differentiation and regulation of apoptosis (8). In SLE the type IFN $\alpha$  can contribute to loss of tolerance and activation of autoreactive T and B cells with production of autoantibodies (9).

The expression of antinuclear antibodies (ANA) can be divided to three stages: antibody induction, maintenance and elimination. Antigens involved in the first two stages may not be the same. Indeed, there is evidence that viral and bacterial antigens can induce ANA production while self molecules may mediate antibody maintenance (10).

However, recent studies by Schroeder et al 2013 (11), Detanico et al 2013 (12) argue that ANA arise

predominantly from nonautoreactive B cells that are transformed into autoreactive cells by the process of somatic hypermutation, which is normally associated with affinity maturation during the germinal center reaction.

CD4 T cell differentiation is altered in SLE presented in the bias of TH1/TH2 balance to TH1 (13) and subsequently transcriptional factors T-bet /GATA3 to T-bet (14), and in the bias of TH17 / Treg to TH17 and subsequently transcriptional factors ROR $\gamma$ /Foxp3 to ROR $\alpha$ t (15,16). Besides the role of follicular helper (TFH) (17) in providing help to B cells allowing the formation of germinal center (18).

CD8 T cell function is impaired in peripheral blood T cells from patients with SLE. Double-negative T cells are also expanded and found within cellular infiltrates in kidney biopsies of patients with lupus nephritis with secretion of IL-1b and IL-17 (19).

### **SLE susceptibility genes**

Here we summarize the confirmed genetic risk loci placed in the context of major SLE disease pathways, where possible.

## **1- Autoantigen generation, clearance & presentation related genes :**

### **1.1- ATG5**

ATG5 which is a regulator of autophagy and apoptosis, was reported to be over-expression in SLE patients that may result in accelerated apoptosis (20) .

### **1.2-GIMAP5**

Several studies have shown that GIMAP5 normally protects cells from apoptosis and that loss of GIMAP5 function results in decreased mitochondrial integrity with lymphocyte apoptosis and lymphopenia, the role of GIMAP5 in the pathogenesis of SLE was confirmed by Hellquist et al 2007(21) .

### **1.3- FcγRs**

Reduced expression of FcγRIIb has been reported for memory B cells and plasma cells from SLE patients. Loss of FcγRIIB results in development of lupus-like symptoms in mice, with the development of autoantibodies and autoimmune glomerulonephritis. On the other hand, reduced function of the FcγRIIIB has been associated with impaired immune complex clearance and hence associated with disease susceptibility (7) .

### **1.4-Complement**

Complement components facilitates clearance of immune complexes and apoptotic cells. So, their deficiency especially in the early components of the classical complement pathway, are associated with SLE. For example, C1q complete deficiency is associated with a 93% risk of SLE . Although related complete deficiencies of C4 or C2 are rare, they as well confer a high risk of developing SLE (22) .

### **1.5- ITGAM**

ITGAM is a receptor for C3b fragment, and thereby shares in the uptake of complement-coated particles and the immune complexes clearance, suggesting that it might be relevant to SLE. Through genome wide association studies (GWASs) in European populations associations between SLE susceptibility and ITGAM have been reported (23,24) .

### **1.6- HLA**

To date, all SLE GWAS in various populations have identified the HLA region as the strongest determinant of genetic risk (25). HLA-DRB1 has been identified as a consistent association locus with SLE in different ethnic populations especially, the

DRB1\*15:01 allele as well as DRB1\*03:01 (22)

## **2- Dendritic cell function and IFN signaling related genes :**

### **2.1- IRF5, IRF7 and IRF8**

IRF5, IRF7 and IRF8 are members of the interferon regulatory factor family. They are a group of transcription factors that play a role in modulation of immune cell growth, differentiation, apoptosis and type I IFN signaling pathway. The associations between SLE risk and polymorphisms of these three gene have been identified in several association studies (26). IRF5 is one of the most strongly associated SLE locus outside the MHC region (27).

### **2.2- TNFAIP3 and TNIP1**

TNFAIP3 and TNIP1 encode regulators of the NF- $\kappa$ B signaling pathway, hence indicating their potential roles in the pathogenesis of SLE. The association between the polymorphisms of TNFAIP3 and SLE were reported through several GWASs with the strongest association at rs2230926 (28).

### **2.3- ETS1 and ELF1**

ETS1 is a transcription factor that binds to IFN stimulated response

elements (ISRE) and function as a negative regulator of the differentiation of Th17 cells and B cells as well as negative regulator of IFN-I induced transcription.(29). ELF1 is a member of ETS family of transcription factors that has roles in T-cell development and function.(30). ETS1 and ELF1 have been identified as a SLE susceptibility loci in Asian populations (27,29).

### **2.4- IFIH1**

IFIH1 is an innate immune receptor for dsRNA that promotes IFN-I production when activated (28). Rs1990760 in IFIH1 has been identified to increase the transcript levels of IFIH1 in SLE patients (31).

### **2.5- TYK2**

Tyrosine kinase 2 (TYK2) is a member of the tyrosine kinase that is important for signaling by type I IFN, also shares in the STAT signaling pathway and induction of Th1 cell differentiation upon antigen stimulation of dendritic cells. Variants in TYK2 with increased expression of type 1 IFN gene were reported to be associated with SLE (32).

### **2.6- UBE2L3**

UBE2L3 is involved in the degra-

dation of Toll-like receptors (TLR) that is widely expressed by lymphocytes<sup>(33)</sup>. GWASs in multiple populations confirmed the association of genetic variations in the UBE2L3 with SLE<sup>(27,31)</sup>

### 2.7- PRKCB

PRKCB is a member of the protein kinase C that is involved in B-cell receptor mediated NF- $\kappa$ B activation<sup>(34)</sup>. A GWAS performed in a Chinese Han population has identified a variant in PRKCB in association with SLE<sup>(35)</sup>.

### 2.8- SLC15A4

SLC15A4 is a member of the solute carrier superfamily of intrinsic membrane transporters. It is involved in Nod1-dependent NF- $\kappa$ B signaling<sup>(36)</sup>. SLC15A4 has been identified as a susceptibility locus for SLE through GWAS in a Chinese population<sup>(27)</sup>

## 3- B cell function and signaling related genes :

### 3.1- BANK1, BLK and LYN

BANK1 encodes for a B-cell-specific scaffold protein involved in B-cell receptor signaling. BLK encodes a non receptor tyrosine kinase that is involved in B-cell receptor sig-

nalizing and B-cell development . Also, LYN encodes a tyrosine protein kinase, which has unique roles in B lymphocyte signaling. The three genes have been reported to be related to SLE<sup>(37)</sup>.

### 3.2- IL-10

IL-10 facilitates B cell proliferation, differentiation and antibody production. The polymorphisms in IL-10 were reported to be associated with SLE in several studies. Furthermore, Increased production of IL-10 by peripheral B cells and monocytes has been shown to correlate with disease activity in SLE patients<sup>(38)</sup>.

### 3.3- RasGRP3

RasGRP3 is a Ras activator expressed in numerous B-cell lines and endothelial cells. A variant of (rs13385731) at the RasGRP3 locus causes suppression of RasGRP3 and is associated with SLE in Chinese Han population<sup>(27)</sup>.

### 3.4- NCF2

Neutrophil cytosolic factor 2 is a critical cytosolic subunit of NADPH oxidase system. It plays a positive role in B-cell activation, resulting in increased autoantibody production<sup>(39)</sup>. In different populations, NCF2

was proven to be associated with increased SLE risk (40) .

### 3.5- PRDM1

PRDM1 is involved in B-cell differentiation and development (27) . Because of the role in B-cell differentiation, the variants that affect PRDM1 could allow the differentiation of plasma cell with more increase in B-cells activity and autoantibody production in SLE patients (41) .

### 3.6- IKZF1

IKZF1 encodes a transcription factor, associated with lymphocyte proliferation and differentiation, and regulation of B-cell receptor signaling(42). Rs4917014 within IKZF1 was identified as a SLE susceptibility locus in a Chinese Han population (27), Reduced expression levels of IKZF1 in peripheral blood are also observed in SLE patients (29) .

## 4- T cells function and signaling related genes :

### 4.1- STAT4

STAT4 is a signaling molecule, in signal transduction by multiple cytokines receptors in T cells and monocytes as IL-17, IL-12, IL-23 and type I IFN, also it mediates the differentiation and proliferation of both T helper

1 (Th1) and Th17 cells which may help the development of autoimmune diseases . Several GWASs have identified STAT4 as a susceptible gene in SLE patients(41) .

### 4.2- PTPN22

PTPN22 ( Protein phosphatase nonreceptor type 22 ) encodes the lymphoid tyrosine phosphate protein, which interacts with cytoplasmic tyrosine kinase and thus inhibits T-cell activation and suppress T regulatory cells. The SNP rs2476601 in PTPN22 was reported to be associated with SLE (43,44) .

### 4.3- TNFSF4

TNFSF4 (also known as OX40L) is a co-stimulatory molecule on the surface of antigen-presenting cells (APCs), that binds to OX40 on activated CD4+and CD8+T cells. Their interaction has been shown in inhibition of the generation of IL-10-producing CD4+type 1 regulatory T cells as well as in the promotion of conventional T cells activation. A risk variant(rs2205960) upstream of TNFSF4 was identified in relation to SLE (45) .

### 4.4- CD44 - PDHX

Two intergenic SNPs located be-

tween PDHX and CD44 on 11p13, (rs2732552 and rs387619), were associated with SLE in a multiethnic study. PDHX encodes a subunit of the pyruvate dehydrogenase complex that is involved in the conversion of pyruvate to acetyl coenzyme A. CD44 encodes a cell-surface glycoprotein, which plays important roles in lymphocyte recirculation, homing, activation and apoptosis (26).

## REFERENCES

1. **Guerra SG, Vyse TJ, Graham DS (2012)** : The genetics of lupus: a functional perspective. *Arthritis Research & Therapy*, 14:211.
2. **Hu S, Wen L, Hu X, et al. (2012)** : IKZF1: a critical role in the pathogenesis of systemic lupus erythematosus?. *Mod Rheumatol*. DOI 10.1007/s10165-012-0706-x.
3. **Janko, C., Schorn, C., Grossmayer, G.E. et al. (2008)** : Inflammatory clearance of apoptotic remnants in systemic lupus erythematosus (SLE). *Autoimmunity reviews.*; 8:9-12.
4. **Munoz, L.E., van Bavel, C., Franz, S. et al. (2008)** : Apoptosis in the pathogenesis of systemic lupus erythematosus. *Lupus.*; 17 : 371-375.
5. **Cohen, P.L. (2006)** : Apoptotic cell death and lupus. *Springer seminars in immunopathology.*; 28:145-152.
6. **Harley, I.T., Kaufman, K.M., Langefeld, C.D., et al. (2009)** : Genetic susceptibility to SLE: new insights from fine mapping and genome-wide association studies. *Nature reviews.*; 10:285-290.
7. **Byrne J, Gabhann J, Elisa Lazari E, et al, . (2012)** : Genetics of SLE: Functional Relevance for Monocytes/Macrophages in Disease .*Clinical and Developmental ImmunologyArticle* .2012; ID 582352.
8. **Ronnblom, L., and Pascual, V.**



- (2008) : The innate immune system in SLE: type I interferons and dendritic cells. *Lupus*. 2008; 17:394-399.
9. **Kaplan, M.H. (2005)** : STAT4: a critical regulator of inflammation in vivo. *Immunologic research*; 31:231-242.
10. **Pisetsky D . (2012)** : The Immunopathogenesis and Immunopathology of Systemic Lupus Erythematosus. In: Schur, P. Massarotti E (eds), *Lupus Erythematosus, Clinical Evaluation and Treatment .*; springer, pp 13-26
11. **Schroeder K, Herrmann M, Winkler. (2013)** : The role of somatic hypermutation in the generation of pathogenic antibodies in SLE . *Autoimmunity.*; Vol. 46, No. 2, Pages 121-127
12. **Detanico T, Clair J, Aviszus K, et al,. (2013)** : Somatic mutagenesis in autoimmunity *Autoimmunity.*; Vol. 46, No. 2, Pages 102-111
13. **Akahoshi M, Nakashima H., Tanaka Y. et al. (1999)** : Th1/Th2 balance of peripheral T helper cells in systemic lupus erythematosus. *Arthritis & Rheumatism.*, Volume 42, Issue 8, pages 1644–1648.
14. **Chan R, Lai F,; Li E et al (2006)** : Imbalance of Th1/Th2 transcription factors in patients with lupus nephritis. *Rheumatology*. 45 : 951 – 957.
15. **Hemdan N., Birkenmeier G., Wichmann G. et al. (2010)** : Interleukin-17-producing T helper cells in autoimmunity. *Autoimmunity Reviews*; 9-11 785-792.
16. Gerli R. Nocentini G, Alunno A et al . (2009) : Identification of regulatory T cells in systemic lupus erythematosus. *Autoimmun Rev.* 8(5) : 426–30.
17. **Nurieva, R, Chung Y, Martinez G et al. (2009)** : Bcl6 mediates the development of T follicular helper

- cells. *Science*. 325, 1001–1005.
18. Dong W, Zhu P, Wang Y, et al . Follicular helper T cells in systemic lupus erythematosus: A potential therapeutic target. *Autoimmunity Reviews*.; 299–304.
19. **Crispin J, Liossis S, Kis-Toth K . et al . (2010)** : Pathogenesis of human systemic lupus erythematosus: recent advances .*Trends in Molecular Medicine*.; Vol. 16 No.2
20. **Yousefi, S., and Simon, H.U. (2007)** : Apoptosis regulation by autophagy gene 5. *Critical reviews in oncology/hematology*.; 63, 241-244.
21. **Hellquist, A., Zucchelli, M., Kivinen, K., Saanalho-Kere, U., et al. (2007)** : The human GIMAP5 gene has a common polyadenylation polymorphism increasing risk to systemic lupus erythematosus. *J Mcd Genet*.; 44:314.321.
22. **Connolly JJ, Hakonarson H. (2012)** : Role of cytokines in systemic lupus erythematosus: recent progress from GWAS and sequencing. *J Biomed Biotechnol*. 798924.
23. **Luo BH, Carman CV, Springer TA. (2007)** : Structural basis of integrin regulation and signaling. *Annu Rev Immunol*.; 25:619-47.
24. **Han S, Kim-Howard X, Deshmukh H, et al. (2009)** : Evaluation of imputation-based association in and around the integrin-alpha-M (ITGAM) gene and replication of robust association between a non-synonymous functional variant within ITGAM and systemic lupus erythematosus (SLE). *Hum Mol Genet*.; 18:1171-80.
25. **Cui Y ,Sheng Y ,Zhang X. (2013)** : Genetic susceptibility to SLE: Recent progress from GWAS, *Journal of Autoimmunity*. ; 1-9.

- 26. Lessard CJ, Adrianto I, Ice JA, et al. (2012)** : Identification of IRF8, TMEM39A, and IKZF3-ZBP2 as susceptibility loci for systemic lupus erythematosus in a large-scale multiracial replication study. *Am J Hum Genet.*; 90:648-60 .
- 27. Han JW, Zheng HF, Cui Y, et al. (2009)** : Genome-wide association study in a Chinese Han population identifies nine new susceptibility loci for systemic lupus erythematosus. *Nat Genet.*; 41:1234-7.
- 28. Flesher DL, Sun X, Behrens TW, et al. (2010)** : Recent advances in the genetics of systemic lupus erythematosus. *Expert Rev Clin Immunol.*; 6:461-79
- 29. Yang W, Shen N, Ye DQ, et al. (2010)** : Genome-wide association study in Asian populations identifies variants in ETS1 and WDFY4 associated with systemic lupus erythematosus. *PLoS Genet.* 2010;6-10.
- 30. Wang D, John SA, Clements JL, et al. (2005)** : Ets-1 deficiency leads to altered B cell differentiation, hyperresponsiveness to TLR9 and autoimmune disease. *Int Immunol.*; 17:1179-91
- 31. Gateva V, Sandling JK, Hom G, et al. (2009)** : A large-scale replication study identifies TNIP1, PRDM1, JAZF1, UHRF1BP1 and IL10 as risk loci for systemic lupus erythematosus. *Nat Genet.*; 41: 1228-33.
- 32. Jarvinen TM, Hellquist A, Koskenmies S, et al. (2010)** : Tyrosine kinase 2 and interferon regulatory factor 5 poly-morphisms are associated with discoid and subacute cutaneous lupus erythematosus. *Exp Dermatol.*; 19:123-31.
- 33. Salloum R, Niewold TB. (2011)** : Interferon regulatory factors in human lupus pathogenesis. *Transl Res.*; 157:326-31.
- 34. Su TT, Guo B, Kawakami Y, et**

- al. (2002)** : PKC-beta controls I kappa B kinase lipid raft recruitment and activation in response to BCR signaling. *Nat Immunol.*; 3:780-6.
- 35. Sheng YJ, Gao JP, Li J, et al. (2011)** : Follow-up study identifies two novel susceptibility loci PRKCB and 8p11.21 for systemic lupus erythematosus. *Rheumatology (Oxford)* ; 50:682-8.
- 36. Lee J, Tattoli I, Wojtal KA, et al. (2009)** : pH-dependent internalization of muramyl peptides from early endosomes enables Nod1 and Nod2 signaling. *J Biol Chem.*; 284:23818-29.
- 37. Ramos PS, Williams AH, Ziegler JT, et al. (2011)** : Genetic analyses of interferon pathway-related genes reveal multiple new loci associated with systemic lupus erythematosus. *Arthritis Rheum.*; 63: 2049-57.
- 38. Deng Y, Tsao BP. (2010)** : Genetic susceptibility to systemic lupus erythematosus in the genomic era. *Nat Rev Rheumatol.*; 6:683-92.
- 39. Vene R, Delfino L, Castellani P, et al. (2010)** : Redoxremodeling allows and controls B-cell activation and differentiation. *Antioxid Redox Signal.*; 13:1145-55.
- 40. Cunninghame Graham DS, Morris DL, Bhangale TR, et al. (2011)** : Association of NCF2, IKZF1, IRF8, IFIH1, and TYK2 with systemic lupus erythematosus. *PLoS Genet*; 7:-10.
- 41. Guerra SG, Vyse TJ, Cunninghame Graham DS. (2012)** : The genetics of lupus: a functional perspective. *Arthritis Res Ther.*; 14:211.
- 42. Yap WH, Yeoh E, Tay A, et al. (2005)** : STAT4 is a target of the hematopoietic zinc-finger transcription factor Ikaros in T cells. *FEBS Lett.*; 579:4470-8.
- 43. Criswell LA, Pfeiffer KA, Lum**

- RF, et al. (2005)** : Analysis of families in the multiple autoimmune disease genetics consortium (MADGC) collection: the PTPN22 620W allele associates with multiple autoimmune phenotypes. *Am J Hum Genet.*; 76:561.
- 44. Harley JB, Alarcon-Riquelme ME, Criswell LA, et al. (2008)** : Genome-wide association scan in women with systemic lupus erythematosus identifies susceptibility variants in ITGAM, PXX, KIAA1542 and other loci. *Nat Genet.*; 40:204-10.
- 45. Cunninghame Graham DS, Graham RR, Manku H, et al. (2008)** : Polymorphism at the TNF superfamily gene TNFSF4 confers susceptibility to systemic lupus erythematosus. *Nat Genet.*; 40: 83-9.